



UAGro
UNIVERSIDAD AUTÓNOMA DE GUERRERO



FACULTAD DE CIENCIAS AGROPECUARIAS Y AMBIENTALES

Influence of Dietary Supplementation of Ensiled Devil Fish and *Staphylococcus saprophyticus* on Equine Fecal Greenhouse Gases Production

TRABAJO DE INVESTIGACIÓN PARA OBTENER EL GRADO DE:
INGENIERO AGRÓNOMO

PRESENTÓ: ERIK DANIEL AGUILAR GARCIA

MATRICULA: 1168225

DIRECTOR DE TESIS: MC. EDSON BRODELI FIGUEROA PACHECO

<https://www.sciencedirect.com/science/article/abs/pii/S073708061930396X#!>

IGUALA DE LA INDEPENDENCIA GRO., JULIO 2019

INDICE

	No. de pag
I. INTRODUCTION	1
II. MATERIALS AND METHODS	2
2.1. Substrate and Treatments	2
2.2. In Vitro Incubation	3
2.3. Calculations and Statistical Analyses	4
III. RESULTS	4
3.1. In Vitro Gas Kinetics	4
3.2. Fecal fermentation parameters	6
IV. DISCUSSION	9
V. CONCLUSIONS	11
VI. REFERENCES	12

INDICE DE TABLAS

Numero de tabla.		No. De Pág.
1	Ingredients and chemical composition of the diets with different levels of ensiled devil fish used as substrates.	3
2	Effect of SS as feed additives on in vitro fecal total gas, CH ₄ , CO ₂ , and H ₂ kinetics of diets at different doses of ensiled DF.	6
3	Effect of SS as feed additives on in vitro fecal fermentation parameters as well as total GP at different incubation periods using different doses of ensiled DF.	7
4	Effect of SS as feed additives on in vitro fecal total gas, CH ₄ , production at different periods using different doses of ensiled DF.	8
5	Effect of SS as feed additives on in vitro fecal CO ₂ , production at different periods using different doses of ensiled DF.	8
6	Effect of SS as feed additives on in vitro fecal H ₂ production at different incubation periods using different doses of ensiled DF.	9

INDICE DE FIGURAS

		No. Pág
Fig1.	Horse fecal total gas, CH ₄ , CO ₂ , and H ₂ production (mL/0.5 g DM) at different incubation periods as affected by the dietary inclusion of ensiled devil fish (DF) at 0 (DF0, control), 6 (DF6), 12 (DF12), and 18% (DF18) of the diet dry matter.	5
Fig. 2	Horse fecal total gas, CH ₄ , CO ₂ , and H ₂ production (mL/0.5 g DM) at different incubation periods as affected by the dietary inclusion of <i>S. saprophyticus</i> (SS) at 0 (SS0, control), 1 (SS1), and 3 mL (SS3) of the diet dry matter.	5

Original Research

Influence of Dietary Supplementation of Ensiled Devil Fish and *Staphylococcus saprophyticus* on Equine Fecal Greenhouse Gases Production

E.D.Aguilar García a, A. Khusro b, E.B.Figueroa Pacheco a, M.J. Adegbeye c, A. Barbabosa-Pliego d, B.Cruz Lagunas a, J.M. Castro Salas a, R. González Mateos a, C. Aarti b, M.M.M.Y. Elghnadour d, *

a) Unidad Académica de Ciencias Agropecuarias y Ambientales, Universidad Autónoma de Guerrero, Iguala de la Independencia, Guerrero, Mexico.

b) Research Department of Plant Biology and Biotechnology, Loyola College, Chennai, Tamil Nadu, India.

c) Department of Animal Science and Livestock Production, Joseph Ayo Babalola University Ikeji-Arakeji Ilesa, Nigeria.

d) Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma del Estado de México, Toluca, México.

ARTICLE INFO

Received 2 May 2019

Received in revised form 24 May 2019

Accepted 31 May 2019

Available online xxx

Keywords:

Devil fish

Staphylococcus saprophyticus

Equine

Greenhouse gases

ABSTRACT

Article history: The present context was designed to investigate the efficacy of devil fish (DF; *Plecostomus* sp.) silage and *Staphylococcus saprophyticus* on fermentation characteristics as well as greenhouse gases production mitigation attributes in horses. Four levels of ensiled DF at 0% (control DF0), 6% (DF6), 12% (DF12), and 18% (DF18) were added into the diet. Moreover, three doses of *S. saprophyticus* (0, 1, and 3 mL/g dry matter [DM]) were used for in vitro fecal

fermentation. The use of ensiled DF resulted in increased ($P < .0001$) pH during fermentation. The asymptotic gas production was the highest ($P < .0001$) in DF6, whereas other supplementation caused lower production than that of control. Lag time for the asymptotic gas production decreased ($P < .05$) with increasing dietary DF doses. Inclusion of *S. saprophyticus* resulted in the lowest ($P < .05$) gas production and mL/0.5 g DM incubated and thus, the reduced gas production up to 23.17% than that of control. The interaction of DF \times *S. saprophyticus* showed the lowest gas production at DF18, whereas the highest production was estimated at DF6 without *S. saprophyticus* after 48 hours. The lowest emission of CO₂ ($P < .0001$) was observed in DF18 inclusion, which was 15.25% lower than that of control at 48 hours of fermentation. In contrast, the lowest hydrogen (H₂) production was estimated in DF0, whereas DF18 exhibited the highest. Inclusion of DF12 and DF18 reduced ($P < .09$) methane (CH₄) emission by 58.24% and 59.33%, respectively. However, DF, *S. saprophyticus*, and DF \times *S. saprophyticus* interaction had no significant effect ($P > .05$) on CH₄ production. In conclusion, ensiled DF and *S. saprophyticus* could be supplemented in equine diet as promising alternatives to corn for mitigating the emission of greenhouse gases effectively

I. INTRODUCTION

Horses are nonruminant, monogastric, and hindgut fermenting herbivores where cecum and colon are the fermentative chambers for disparate microbiota. The microbial population present in hindgut are known for the stimulation of immunity, exclusion of pathogens, and detoxification of hazardous components [1]. The diet of horses is enriched with fibers; however, hindgut microbiota enables digesting fiber-based diets gradually due to the fact that the fiber is indigestible by secreted enzymes. However, alteration in feeding practices and activities of modern-day horses have led to an increased level of grain or starch and lowered levels of fiber in their diet [2,3]. This is done to provide quick energy release to meet the energy need of high-paced activity of equine [4]. However, such feeding practices lead to the leading causes of several disorders, namely gastric ulceration, hindgut acidosis, and endotoxemia [5]. In addition, feeding such diets may decrease the starch digestion trait in the small intestine and alter the microbial population as well as fibrolytic characteristics in the hindgut, thereby reducing the ability to use energy from the diets, as a result of alteration in hindgut pH [4,6]. However, pectin-rich by-product (lemon, tangerine, and pineapple) and agro-based by-product (sugar beet pulp and soybean hull) have been put forward to provide energy for horses without causing digestive disturbances or offset [4]. Still, there is an urgency to explore other resources for providing the energy demands, intestinal health, and enhance the athletic high-level performances of modern horses.

Currently, the supplementation of diversified additives into the feeds is considered auspicious strategies to enhance the energy utilization in horses. Unfortunately, the perpetual emission of greenhouse gases (GHG), particularly methane (CH₄) and carbon dioxide (CO₂), from animals due to the fermentation is the colossal burden globally. These GHGs are considered not only environmental pollutants but also hazardous to human health, resulting in global warming [7]. The quest for auspicious natural alternative resources to mitigate the emission of GHG for cleaner society and sustainable environment has gained immense interest. For instance, distinct natural feed additives such as plant extract [8], enzyme [9], yeasts [7], and lactobacilli [10] had been used. Nevertheless, the exploitation of coagulase-negative staphylococci (CNS) as feed additive in horse nutrition for mitigating the emission of GHG is not evidenced yet. *Staphylococcus equorum*, *S. hominis*, *S. cohnii*, *S. capitis*, *S. condimentii*, *S. succinus*, and *S. xylosus* belong to CNS group [11]. In recent times, CNS have emerged as the prevalent heterogeneous group of bacteria and included under Qualified Presumption of Safety status by the European Food Safety Authority Scientific Committee on a case-by-case basis within a particular taxonomic group [12].

In recent years, CNS have emerged as different group of fermented food-associated bacteria revealing probiotic properties [13]. In addition, CNS were reported as the predominant type of bacteria in some Korean fermented food [14]. This is an indication of the fermenting properties of CNS or its probiotic properties. Furthermore, devil fish (DF) (*Plecostomus* sp.) are included in animal's diet because of its abundance and maximum digestibility [15]. In addition, ensiling the fish could pave a way of improving its usage as feed ingredient. The proteolytic enzyme in the ensiled fish could improve feed digestibility. Besides, it is a well-established fact that fermented foods are enriched with health beneficial probiotic microbes [16]. The effect of DF has been studied in ruminant diet [17] with a better response in fermentation kinetics. However, this kind of investigation is unexplored in equine.

Considering this, a further significant attempt was undertaken in this context to fill the gap of research by determining the fermentation kinetics and GHG production mitigation attributes of *S. saprophyticus* and DF in horses as ideal alternatives to feed supplements for a cleaner and ecofriendly product.

Animal welfare/ethical statement: The research was performed in accordance with the ethical standard laid down in the 1996 Declaration of Helsinki and its later amendments.

Conflict of interest statement: The authors declare no conflicts of interest.

* Corresponding author at: M.M.M.Y. Elghnadour, Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma del Estado de México, Toluca 50000, Mexico. *Email address:* mmohamede@uaemex.mx (M.M.M.Y. Elghnadour)

<https://doi.org/10.1016/j.jevs.2019.05.023 0737-0806/>

II. MATERIALS AND METHODS

2.1. Substrate and Treatments

Substrate (diets) used in this study were dried at 60°C for 48 hours before in vitro incubation. The diet level was 0%, 6%, 12%, and 18% of ensiled DF of diet dry matter (DM) and represented as DF0, DF6, DF12, and DF18, respectively. In addition, three doses (0, 1, and 3 mL/g DM substrate) of *S. saprophyticus* (SS) at 5×10^{11} CFU/g represented as SS0, SS1, and SS3 were used for in vitro fermentation. Diet formulation and chemical composition of diets are shown in

Table 1.

To ensile the DF, the fresh live fish was obtained from the Tuxpan lagoon municipality of the city of Iguala Gro. The fish was washed with water to remove

the soil and particles stuck to the fish. After milling process, 5 kg of fish were mixed with 14 L of molasses and 1 L of natural yogurt in a bucket with a capacity of 20 L in which an airtight lid was placed to avoid leaks and air entrances, and then it was kept for 30 days. The cuvette was opened after 30 days and mixed with the ingredients as mentioned in Table 1.

2.2. In Vitro Incubation

Horses were fed the compounded diet (substrate) ad libitum and provided fresh water for 7 days before collection phase. Fecal content (inoculum source) collected from the rectum were obtained from four Azteca horses (aged 5–8 years, 480 ± 20.1 kg). Culture broth

Table 1

Ingredients and chemical composition of the diets with different levels of ensiled devil fish used as substrates.^a

Ingredients	DF0	DF6	DF12	DF18
Ground corn	73.5	67.5	61.5	55.5
Pastures	15	15	15	15
Soybean meal	9	9	9	9
Devil fish	0	6	12	18
Minerals	2.5	2.5	2.5	2.5
Chemical composition (%)				
Organic matter	3.4	3.5	4.01	4.10
Ether extract	11.4	12.2	11.3	12.13
Acid detergent fiber	2.7	2.8	2.7	3.1
Crude protein	45.3	45.5	46.2	47.9
Neutral detergent fiber	4.7	4.9	4.89	5.02
Acid detergent lignin	2.1	2.3	2.1	2.3
Minerals ^b	34.7	34.7	34.7	34.7

^a Adapted from Abrego Salgado [18].

^b Cu: 21.18 ppm, Fe: 4971.66 ppm, Zn: 343.75 ppm, Ca: 9.96%, Mg: 0.2495%, K: 0.8895%, Na: 1.296%, Pb: 0.0029%, P: 14.395%, S: 3.125%.

was added to the fecal contents in a ratio of 4:1 and kept under CO₂ environment throughout the entire in vitro incubation process (39°C; 48 hours). All incubations were performed in triplicate, and either rumen fluid or fecal fluid was used as a blank. Data at 2, 4, 6, 8, 10, 12, 14, 24, and 48 hours using the pressure reading technique was used to estimate total gas, CO₂, CH₄, and CO₂ emissions [19]. CO₂, CH₄, and H₂ concentrations were also measured in the headspace of the bottles up to 48 hours using the gas detector (AIR QUALITY MONITOR YesAIR, Critical Environment Technologies Canada Inc, Delta, British Columbia, Canada). Furthermore, pH was measured, and DM degradability (DMD) was estimated after filtration [20].

2.3. Calculations and Statistical Analyses

Kinetic parameters of gas production (mL/g DM) were calculated according to France et al. [21] using the NLIN option of SAS [22]. The DMD was calculated according to the methodology of Menke et al [23]. Fecal fermentation data were estimated as a completely randomized design as per PROC GLM option:



$$Y_{ij} = \mu + B_i + \varepsilon_{ij}$$

where, Y_{ij} = observation obtained with i th level of LAB;
 B_i = level of LAB ($i = 1-4$); μ = general mean; ε_{ij} = experimental error. Linear and quadratic polynomial contrasts were implied to assess responses for increasing concentrations of *S. saprophyticus*. Turkey's test was used to calculate multiple comparisons among means. Significance level was estimated at $P < .05$.

III. RESULTS

3.1. In Vitro Gas Kinetics

Figs. 1 and 2 showed the effect of ensiled DF and *S. saprophyticus* on horse fecal total gas, CH₄, CO₂, and H₂ production. Inclusion of *S. saprophyticus* had no significant effect ($P > .05$) on total gas, CH₄, CO₂, and H₂ production. Furthermore, Table 2 showed that ensiled DF had a linear effect ($P < .0001$) on the asymptotic gas production ($P = .031$), rate of gas production, and lag time ($P < .0001$). The DF6 showed the highest asymptomatic gas production and CO₂ production, whereas the DF18 exhibited the lowest gas production and CO₂ emission. Interaction of DF \times *S. saprophyticus* had no effect ($P > 0.05$) on CH₄, CO₂, and H₂ except for the asymptotic gas production ($P = 0.0017$) and the lag time ($P = 0.039$).






Fig. 1: Horse fecal total gas, CH₄, CO₂, and H₂ production (mL/0.5 g DM) at different incubation periods as affected by the dietary inclusion of ensiled devil fish (DF) at 0 (DF0, control), 6 (DF6), 12 (DF12), and 18% (DF18) of the diet dry matter.




Fig. 2: Horse fecal total gas, CH₄, CO₂, and H₂ production (mL/0.5 g DM) at different incubation periods as affected by the dietary inclusion of *S. saprophyticus* (SS) at 0 (SS0, control), 1 (SS1), and 3 mL (SS3) of the diet dry matter.

3.2. Fecal fermentation parameters

The devil fish supplementation showed a linear effect ($P < 0.0001$) on the pH and gas production at 24 and 48 h. The DF12 and DF0 showed the highest and the lowest pH values, respectively. Furthermore, DF6 and DF18 doses resulted in the highest and the lowest mL gas/0.5 g DM at 24 and 48 h. Similarly, *S. saprophyticus* had a linear effect ($P < 0.05$) on gas production but it had no effect ($P > 0.05$) on fecal pH. There was a linear decrease ($P < 0.05$) in gas production in a dose dependent manner at 24 and 48 h. Devil fish \times *S. saprophyticus* estimated linear effect ($P < 0.002$) on gas production (mL/0.5 g DM incubated) at 48 h of incubation. The devil fish, *S. saprophyticus*, and devil fish \times *S. saprophyticus* interaction showed no significant effect on DMD. Similarly, *S. saprophyticus* exhibited no influence on the rumen fluid pH (Table 3).

Table 2. Effect of SS as feed additives on in vitro fecal total gas, CH₄, CO₂, and H₂ kinetics^a of diets at different doses of ensiled DF.

DF Doses	SS Doses	Total gas			CH ₄			CO ₂			H ₂		
		<i>b</i>	<i>c</i>	<i>Lag</i>	<i>b</i>	<i>c</i>	<i>Lag</i>	<i>b</i>	<i>c</i>	<i>Lag</i>	<i>b</i>	<i>c</i>	<i>Lag</i>
DF0	0	128.6	0.008	4.439	10.4	0.001	5.908	89.7	0.004	4.409		0.011	6.750
	1	132.8	0.003	6.042	11.9	0.001	6.941	78.1	0.001	4.844		0.007	6.695
	3	128.5	0.010	4.989	12.7	0.001	7.662	90.2	0.001	4.692		0.027	6.913
DF6	0	143.8	0.009	3.764	20.4	0.023	7.990	88.6	0.011	4.265		0.007	6.206
	1	132.8	0.005	4.370	13.7	0.001	7.761	91.8	0.012	2.957		0.020	5.815
	3	135.8	0.005	1.965	18.0	0.020	8.098	81.2	0.004	3.800		0.013	5.279
DF12	0	136.3	0.003	2.072	16.3	0.001	6.882	88.0	0.009	4.278		0.016	9.479
	1	133.3	0.004	6.276	4.8	0.001	7.560	97.1	0.006	3.643		0.013	5.712
	3	91.8	0.009	1.724	3.0	0.002	8.837	77.1	0.003	3.930		0.007	6.859
DF18	0	88.9	0.001	1.753	7.5	0.011	8.220	60.2	0.012	4.460		0.020	6.848
	1	98.9	0.004	1.632	7.6	0.001	5.079	62.8	0.010	3.988		0.018	6.070
	3	105.3	0.002	1.917	28.4	0.005	9.203	62.0	0.005	3.087		0.009	6.416
<i>P</i> values													
DF													
Linear		<.0001	.0307	<.0001	.8301	.446	.3764	<.0001	.0307	.2095		.8764	.6646
Quadratic		.1532	.6781	.8436	.6575	.7236	.3636	.0012	.9366	.5882		.4613	.2863
SS													
Linear		.0557	.4179	.5026	.3539	.6908	.0721	.2886	.0455	.3868		.9329	.1695
Quadratic		.252	.2624	.0008	.2567	.1087	.0787	.3884	.6551	.5872		.8841	.1988
DF \times SS		.0017	.3773	.0392	.3405	.6714	.2263	.1323	.9699	.8475		.2551	.5215

Abbreviations: DF, devil fish; GP, gas production; SS, *S. saprophyticus*.

^a *b* is the asymptotic GP (mL/g DM); *c* is the rate of GP (per hour); *Lag* is the initial delay before GP begins (hour).

3.3. Fecal greenhouse gas production

Table 4 showed that devil fish doses, *S. saprophyticus* concentrations, and devil fish \times *S. saprophyticus* interaction had neither linear nor quadratic effect ($P > 0.05$) on mL CH₄/0.5g DM incubated, mL CH₄/0.5 g DM degraded, and proportional CH₄ production. At 8 h of incubation period, there was complete absence of in vitro fecal CH₄ production. However,

devil fish and *S. saprophyticus* doses revealed quantitative reduction in CH₄ production (Fig. 2).

Table 5 showed that devil fish doses had linear effect ($P < 0.05$) on mL CO₂/0.5 g DM incubated and mL CO₂/0.5 g DM degraded but showed no effect on the proportional CO₂ production. The DF6 and DF12 revealed the highest mL CO₂/0.5g DM degraded and mL CO₂/0.5g DM incubated while DF18 exhibited the lowest mL CO₂/0.5g DM degraded and mL CO₂/0.5g DM incubated at 24 and 48 h of incubation. Furthermore, devil fish × *S. saprophyticus* interaction had a linear effect ($P < 0.05$) on the mL CO₂/0.5 g DM degraded and proportional CO₂ production with DF6 exhibiting the highest while DF18 revealing the lowest CO₂ production. However, *S. saprophyticus* estimated no significant ($P > 0.05$) impact on CO₂ emission (Fig. 2).

Table 6 showed that devil fish doses had a linear ($P < 0.02$) influence on the proportional H₂ production. The DF18 produced the highest H₂, while DF0 quantified the lowest H₂ production. *S. saprophyticus* doses and devil fish × *S. saprophyticus* interaction had no significant ($P > 0.05$) impact on mL H₂/0.5 DM incubated and mL H₂/0.5 g DM degraded. In contrary, *S. saprophyticus* doses and devil fish × *S. saprophyticus* interaction exhibited linear effect ($P < 0.03$) on mL H₂/0.5 DM incubated at 12 h (Fig. 2).

Table 3

Effect of SS as feed additives on in vitro fecal fermentation parameters as well as total GP at different incubation periods using different doses of ensiled DF.

DF Doses	SS Doses	Fermentation Parameters		GP mL/0.5 g DM Incubated				GP mL/0.5 g DM Degraded			
		pH	DMD	8	12	24	48	8	12	24	48
DF0	0	6.3	73.4	36.7	48.1	78.3	129.9	26.9	35.2	57.5	95.5
	1	6.3	76.8	32.3	41.4	72.9	131.9	24.9	32.0	56.2	101.5
	3	6.4	74.8	30.3	40.7	72.7	135.7	22.7	30.5	54.4	101.5
DF6	0	6.6	77.5	37.0	49.1	86.6	148.7	28.6	38.0	67.1	115.3
	1	6.5	77.5	34.8	47.1	79.3	136.2	27.0	36.5	61.5	105.6
	3	6.6	76.2	33.8	43.4	73.8	136.3	25.8	33.0	56.2	103.8
DF12	0	6.6	76.8	36.4	46.8	79.3	137.5	27.9	35.9	60.8	105.5
	1	6.6	78.7	32.7	42.4	72.2	134.5	25.7	33.4	56.8	105.9
	3	6.6	75.4	27.4	35.0	58.0	102.6	20.7	26.4	43.8	77.4
DF18	0	6.6	78.2	30.1	40.2	57.7	99.8	23.5	31.4	45.1	78.0
	1	6.5	66.7	32.0	38.8	57.6	100.6	21.4	26.3	38.7	67.8
	3	6.5	72.9	31.8	42.7	56.9	106.4	23.2	31.1	41.8	77.8
<i>P</i> values											
DF											
Linear		<.0001	.4416	.2547	.1813	<.0001	<.0001	.1842	.1733	.0001	<.0001
Quadratic		<.0001	.2469	.9638	.7489	.1128	.0167	.477	.6607	.0798	.0333
SS											
Linear		.2386	.5432	.0046	.0041	.0002	.0061	.0127	.0135	.0037	.0507
Quadratic		.038	.7678	.9985	.6017	.9759	.6453	.9034	.6828	.9899	.8136
DF × SS		.3319	.554	.1949	.1628	.0978	.0002	.6221	.4442	.3747	.0345

Abbreviations: DF, devil fish; DMD, dry matter degradability (%); GP, gas production; SS, *S. saprophyticus*.

Table 4
Effect of SS as feed additives on in vitro fecal CH₄ production at different incubation periods^a using different doses of ensiled DF.

DF Doses	SS Doses	mL CH ₄ /0.5 g Dry Matter Incubated			mL CH ₄ /0.5 g Dry Matter Degraded			Proportional CH ₄ Production		
		12	24	48	12	24	48	12	24	48
DF0	0	0.8	3.2	11.8	0.6	2.4	8.7	1.7	4.1	9.1
	1	1.3	4.0	12.8	1.0	3.1	9.9	3.0	5.4	9.8
	3	0.6	3.2	14.1	0.4	2.4	10.7	1.4	4.4	10.4
DF6	0	0.8	4.5	19.0	0.6	3.5	14.8	1.5	5.2	12.5
	1	0.4	3.2	14.5	0.3	2.5	11.3	0.8	4.0	10.4
	3	1.0	4.6	19.1	0.7	3.5	14.5	2.2	6.1	13.9
DF12	0	0.7	4.2	17.2	0.6	3.2	13.2	1.6	5.2	12.6
	1	0.6	1.8	6.1	0.5	1.4	4.8	1.4	2.5	4.5
	3	0.5	1.2	3.9	0.3	0.9	2.9	1.2	2.0	3.8
DF18	0	0.9	2.4	8.1	0.7	1.9	6.3	2.1	4.2	8.0
	1	0.5	2.1	8.5	0.4	1.5	6.2	1.3	3.6	8.3
	3	0.3	2.3	11.3	0.2	1.6	8.1	0.7	4.1	10.7
<i>P</i> values										
DF		.53	1.39	5.25	.42	1.11	4.13	1.14	1.68	3.71
Linear		.2246	.0811	.1557	.2199	.0839	.1498	.2512	.4026	.6746
Quadratic		.5725	.3876	.3511	.6307	.4802	.4475	.5462	.1393	.1267
SS										
Linear		.3329	.1848	.3794	.3241	.1695	.3198	.4788	.4373	.5902
Quadratic		.9467	.4069	.1772	.9143	.471	.2167	.8733	.36	.1645
DF × SS		.4385	.3163	.1378	.4365	.3615	.1702	.3139	.2499	.1515

Abbreviations: DF, devil fish; DMD, dry matter degradability (%); GP, gas production; SS, *S. saprophyticus*.¹

^a No detect CH₄ production before 12 hr of incubation.

Table 5
Effect of SS as feed additives on in vitro fecal CO₂ production at different incubation periods using different doses of ensiled DF.

DF Doses	SS Doses	mL CO ₂ /0.5 g Dry Matter Incubated				mL CO ₂ /0.5 g Dry Matter Degraded				Proportional CO ₂ Production			
		8	12	24	48	8	12	24	48	8	12	24	48
DF0	0	6.9	21.7	39.5	90.4	5.1	15.8	29.0	66.3	18.7	44.7	50.2	69.5
	1	5.3	16.1	34.5	79.1	4.0	12.4	26.4	60.4	17.0	39.7	48.2	60.3
	3	6.5	18.4	39.4	91.4	4.9	13.8	29.5	68.3	21.7	45.3	54.2	67.4
DF6	0	8.1	22.2	45.5	89.5	6.2	17.2	35.2	69.3	21.7	45.3	52.5	60.5
	1	10.1	26.1	48.8	92.9	7.8	20.2	37.8	72.0	29.0	55.3	61.6	68.4
	3	6.1	20.2	37.5	82.6	4.6	15.4	28.6	62.9	18.0	46.7	50.8	60.6
DF12	0	6.8	22.2	41.9	89.2	5.2	17.0	32.2	68.5	18.3	47.3	52.9	65.0
	1	8.1	24.4	46.6	98.1	6.3	19.2	36.8	77.4	24.7	57.7	64.6	72.9
	3	5.8	17.9	35.0	77.8	4.4	13.5	26.4	58.7	21.3	51.7	60.8	76.1
DF18	0	5.3	16.4	28.8	60.8	4.2	12.8	22.5	47.6	17.7	40.9	49.9	61.0
	1	5.3	16.1	31.7	63.7	3.6	10.9	21.4	42.1	16.7	41.9	55.5	63.7
	3	5.4	17.9	27.9	62.7	3.9	13.1	20.5	46.0	17.0	42.0	49.2	58.9
<i>P</i> values													
DF													
Linear		.2025	.1346	.0014	<.0001	.1500	.1055	.0029	<.0001	.2771	.5155	.8331	.2309
Quadratic		.0759	.0021	.0009	.001	.0269	.0009	.0007	.0005	.0427	.0001	.0072	.0217
SS													
Linear		.1707	.0745	.0598	.3033	.1237	.0643	.0656	.2301	.7914	.3955	.4048	.5968
Quadratic		.117	.2578	.0556	.3705	.1116	.278	.1015	.4818	.0717	.1048	.0552	.6017
DF × SS		.0799	.0162	.1032	.1218	.054	.028	.1714	.1397	.0412	.1462	.382	.3221

Abbreviations: DF, devil fish; DMD, dry matter degradability (%); GP, gas production; SS, *S. saprophyticus*.

Table 6Effect of SS as feed additives on in vitro fecal H₂ production at different incubation periods using different doses of ensiled DF.

DF Doses	SS Doses	mL H ₂ /0.5 Dry Matter Incubated				mL H ₂ /0.5 g Dry Matter Degraded				Proportional H ₂ Production					
		8	12	24	48	8	12	24	48	8	12	24	48		
DF0	0	0.11	4.22	7.77	16.89	22.60	0.09	3.14	5.78	12.55	17.61	0.33	10.0	13.00	17.00
	1	0.89	4.67	9.03	23.99	29.48	0.69	3.66	7.07	17.97	22.87	2.67	12.0	17.67	19.67
	3	0.75	4.34	8.48	23.96	32.17	0.56	3.26	6.37	18.59	24.52	2.67	11.6	17.67	23.67
DF6	0	1.24	5.91	11.85	26.69	22.02	0.96	4.58	9.19	20.45	3.33	13.6	13.6	19.33	16.33
	1	0.35	5.63	10.81	14.86	25.68	0.27	4.37	8.39	17.35	1.00	19.6	14.33	25.67	
	3	0.34	6.47	14.38	25.41	0.26	4.93	10.96	11.21	1.00	1.00	14.6	25.00		
DF12	0	0.36	5.56	11.79	24.52	0.28	4.25	9.02	20.08	17.43	1.00	11.3	23.00		
	1	0.33	3.96	8.20		0.26	3.10	6.44	17.89	1.00	10.6	21.6			
	3	0.27	2.78	6.30		0.21	2.10	4.76		1.00	21.0				
DF18	0	0.30	6.56	12.49	12.29	.3036	0.24	5.12	9.76		1.00	18.6	.0018		
	1	0.32	7.17	10.76		.5556	0.21	5.01	8.50	.4507	1.00		.1252		
	3	0.32	6.06				0.23	4.34	7.88	.7401	1.00				
<i>P</i> values					.8118								.0045	.9422	
DF				.089	.7869				.6949			.0485	.7695	.6997	
Linear		.1937	.0279	.4194	.4525	.1737	.0702	.1574	.7697	.1701					
Quadratic		.4763	.0956			.5171	.1751	.5543	.5049	.4222		.8423			
SS				.5578								.774			
Linear		.6329	.429	.7878		.5744	.3693	.4958		.895		.6734			
Quadratic		.9496	.8646	.6738		.9532	.9041	.7604		1.000					
DF × SS		.0589	.787			.0599	.8652	.7372		.099					

Abbreviations: DF, devil fish; DMD, dry matter degradability (%); GP, gas production; SS, *S. saprophyticus*.

IV. DISCUSSION

At present, improving the nutrition through the use of unconventional ingredients of crops or animal sources, novel additives, and microorganisms are the most common practises of researchers and farmers. In addition, reducing greenhouse gases productions are also an important factor in equine nutritional interventions. Total gas production is an indication of feed digestibility or degradation. In this context, total gas production was estimated to be increased due to the supplementation of devil fish. The increment in the total gas production by DF6 with respect to the control may be attributed to the enzymatic activities such as protease and lipase from the devil fish, which enhanced the growth of microbes. It could also be due to the fact that the high protein content leads to more availability of ammonia nitrogen, which enhanced their growth. Makkar et al [23] reported that protein fermentation produced lesser gas compared to carbohydrate. Hence, in contrary to our study, the highest protein supplementation resulted in the reduced gas production. Velazquez et al [24] stated that Lag time is a measure of the time required for feed digestibility by gut microbes to initiate digestibility. In this investigation, Lag time was reduced during ensiling due to the fermentation process. The lower Lag time with devil fish supplementation may be attributed to the ability of microbes to adapt or reveal probiotic traits [16] Therefore, it could be stated that devil fish had some probiotics properties, which led to quick adherence and colonization of feed particles compared to the control.

In this study, total CO₂ emissions from horses were reduced in the presence of high doses of devil fish. The highest total CO₂ production in DF12 and its similarity with DF6 could be due to the fermentation process. However, the lowest total CO₂ in DF18 may be attributed to the high crude protein content of devil fish. The similar pattern was observed in mL CO₂/0.5 g DM incubated and degraded too (Table 5). Velázquez et al [25] demonstrated that the lower CO₂ production may be influenced by the high protein content in a diet. Additionally, the ammonia–N nitrogen accumulation in the medium might have prevented the release of CO₂ in the bottle [26].

Faniyi et al [27] reported that pH is a fermentation parameter that quantifies the state of acidity and alkalinity in the gut and during fermentation. Similarly, the characteristics of a feed consumed by animal influence the pH. In another words, during in vitro assay, the fluid pH is influenced by the substrate characteristics. In the present investigation, pH value was increased due to the supplementation of varied doses of devil fish. The increase in pH with devil fish supplementation may be attributed to the high-protein and low-carbohydrate in the substrate fermented, compared to the control, which had higher ground corn in it [28,29].

Elghandour et al [30] reported that H₂ removal stimulates bacteria during digestion. This indicates that the higher proportional H₂ in DF18 throughout incubation period might have affected digestion. In contrast, the higher H₂ gas in DF6 could be an indication of production of more acetate and butyrate where H₂ is produced in the process [31]. The numerical reduction in CH₄ production at 24 h of incubation by DF12 and DF18 may be attributed to the lower production of CO₂ and H₂ gas.

Borah et al [16] had reported the antagonistic activity of some species of *Staphylococcus*. In the present study, we observed lower gas production (mL/0.5 g DM incubated and degraded) with increasing doses of *S. saprophyticus*. This indicates that *S. saprophyticus* had some inherent antimicrobial and growth inhibitory properties. Besides, Khusro et al [11] reported that *Staphylococcus* sp. showed lack of amylolytic activity, which exhibited their inability to degrade starch. Therefore, the lower gas production encountered with *S. saprophyticus* supplementation may be attributed to the fact that bacteria are unable to degrade starch. Furthermore, Laukova´ and Marekova´ [32], Sung et al [33], and Khusro et al [13] had reported that CNS strains are ideal producers of bacteriocins. Thus, the bacteriocin produced by *S. saprophyticus* might have inhibited the fermentative microorganisms. The supplementation of *S. saprophyticus* at distinct doses revealed reduction in CH₄ emission. Low CH₄ production with *S. saprophyticus* supplementation may be explained by their ability to reduce nitrate to nitrite [34,35]. Furthermore, the present study showed that *S. saprophyticus* supplementation had no significant impact on CO₂ and H₂ emission. The presence of staphylococci

might have enhanced formation which would serve as terminal electron acceptors during fermentation of feeds [36].

Over the past few years, a significant attempt had been undertaken not only to improve the nutritional quality but also to reduce the emission of greenhouse gases from livestock through the synergistic role of additives. In this regard, synergistic role of fibrous forages along with live yeasts and fibrolytic enzymes have been successfully investigated towards the improved fermentation as well as mitigation of CH₄ and CO₂ emission from equine [37-39]. The present investigation further filled the gap of equine research by evaluating the pivotal synergistic role of devil fish and *S. saprophyticus* as promising feed additives in the reduction of greenhouse gases emission from horses.

V. CONCLUSIONS

Inclusion of devil fish at 6% of diet can improve feed gas production, without disrupting the gut pH. The supplementation of DF12 and DF18 reduced CH₄ production by 58.24 and 59.33%, respectively. DF18 reduced total CO₂ production by 15.25%, while SS1 and SS3 mitigated CH₄ production by 50.54 and 58.24%, respectively. Similarly, the low gas production pattern with *S. saprophyticus* supplementation indicates its antimicrobial properties, suggesting good prospect in livestock nutrition. The devil fish and *S. saprophyticus* could be potential feed additives as alternatives to the conventional antibiotics.

VI. REFERENCES

1. Shepherd ML, Swecke Jr WS, Jr Jensen RV, Ponder MA. Characterization of the fecal bacteria communities of forage-fed horses by pyrosequencing of 16S rRNA V4 gene amplicons. *FEMS Microbiol Lett* 2012;326:62–8
2. de Fombelle A, Varloud M, Goachet AG, Jacotot E, Philippeau C, Drogoul C, Julliand V. Characterisation of the microbial and biochemical profile of the different segments of the digestive tract in horses fed two distinct diets. *Anim Sci* 2003;77:293-304.
3. Sadet-Bourgeteau S, Julliand V. Equine microbial gastro-intestinal health in: the impact of nutrition on the health and welfare of horses. In: Presented at the 5th European Workshop equine nutrition, EAAP Scientific Series, Cirencester, United Kingdom; 2010; p. 161-182.
4. Cipriano-Salazar M, Adegbeye MJ, Elghandour MMY, Barbabosa-Pilego A, Mellado M, Hassan A, Salem AZM. The dietary components and feeding management as options to offset digestive disturbances in horses. *J Equine Vet Sci* 2019;74:103-10
5. Rowe JB, Lees MJ, Pethick DW. Prevention of acidosis and laminitis associated with grain feeding in horses. *J Nutr* 1994;124:2742-4.
6. Mungall BA, Kyaw-Tanner M, Pollitt CC. *In vitro* evidence for a bacterial pathogenesis of equine laminitis. *Vet Microbiol* 2001;79:209–23.
7. Elghandour MMY, Vazquez JC, Salem AZM, Kholif AE, Cipriano MM, Camacho LM, Marquez O. *In vitro* gas and methane production of two mixed rations influenced by three different cultures of *Saccharomyces cerevisiae*. *J Appl Anim Res* 2017;45:389-95.

8. Cedillo J, Kholif AE, Salem AZM, Elghandour MMY, Vazquez JF, Alonso MU, Barbabosa A, Chagoyan JCV, Reyna AG. Oral administration of Sauce Iloron extract to growing lambs to control gastrointestinal nematodes and *Moniezia* spp. *Asian Pac J Trop Med* 2015;8:520-5.
9. Vallejo LH, Salem AZM, Kholif AE, Elghangour MMY, Fajardo RC, Rivero N, Bastida AZ, Mariezcurrena MD. Influence of cellulase or xylanase on the *in vitro* rumen gas production and fermentation of corn stover. *Indian J Anim Sci* 2016;86:70-4.
10. Elghandour MMY, Khusro A, Greiner R, Salem AZM, Lugo de la Fuente J, Marquez-Molina O, Barbabosa-Pilego A, Montes-de-Oca Jimenez R. Horse fecal methane and carbon dioxide production and fermentation kinetics influenced by *Lactobacillus farciminish* supplemented diet. *J Equine Vet Sci* 2018;62:98-101.
11. Khusro A, Aarti C, Barbabosa-Pilego A, Hernández SR. Anti-pathogenic, antibiofilm, and technological properties of fermented food associated *Staphylococcus succinus* strain AAS2. *Prep Biochem Biotech* 2019;49:176-83
12. EFSA. Introduction of a Qualified Presumption of Safety (QPS) approach for assessment of selected microorganisms referred to EFSA1. Opinion of the Scientific Committee (Question No EFSA-Q-2005-293. *EFSA J.* 2007;587:1–16. Retrieved from www.efsa.europa.eu/en/scdocs/doc/587.pdf
13. Khusro A, Aarti C, Salem AZM, Rodríguez GB, Rivas-Cáceres RR. Antagonistic trait of *Staphylococcus succinus* strain AAS2 against uropathogens and assessment of its *in vitro* probiotic characteristics. *Microb Pathog* 2018;118:126–32
14. Guan L, Cho KH, Lee JH. Analysis of the cultivable bacterial community in jeotgal, a Korean salted and fermented seafood, and identification of its dominant bacteria. *Food Microbiol* 2011;28:101–13.

15. Avdalov N, Barlocoo N, Bauza R, Giacommeti L, Panucio A. Evaluación del ensilaje biológico de pescado en la alimentación de cerdos en engorde. Instituto Nacional de Pesca. Segunda consulta de expertos sobre tecnología de productos pesqueros en América latina. Montevideo, Uruguay. 1989; pp. 88-98.
16. Borah D, Gogoi O, Adhikari C, Kakoti BB. Isolation and characterization of the new indigenous *Staphylococcus* sp. DBOCP06 as a probiotic bacterium from traditionally fermented fish and meat products of Assam state. Egypt J Basic Appl Sci 2016; 3:232–40.
17. Tejeda-Arroyo E, Cipriano-Salazar M, Camacho-Díaz LM, Salem AZM, Kholif AE, Elghandour MMY, DiLorenzo N, Cruz-Lagunas B. Diet inclusion of devil fish (*Plecostomus* spp.) silage and its impacts on ruminal fermentation and growth performance of growing lambs in hot regions of Mexico. Trop Anim Health Prod 2015;47:861–6.
18. Theodorou MK, Williams BA, Dhanoa MS, McAllan AB, France J. A simple gas production method using a pressure transducer to determine the fermentation kinetics of ruminant feeds. Anim Feed Sci Tech 1994;48:185-97.
19. Elghandour MM, Chagoyan JCV, Salem AZM, Kholif AE, Castañeda JSM, Camacho LM, Cerrillo-Soto MA. Effects of *Saccharomyces cerevisiae* at direct addition or pre incubation on *in vitro* gas production kinetics and degradability of four fibrous feeds. Ital J Anim Sci 2014;13:295-301.
20. France J, Dijkstra J, Dhanoa MS, Lopez S, Bannink A. Estimating the extent of degradation of ruminant feeds from a description of their gas production profiles observed *in vitro*: derivation of models and other mathematical considerations. Br J Nutr 2000;83:143-50.

21. SAS. Statistical Analysis System. User's Guide: Statistics. Ver 9.0. SAS Institute, Cary, NC. 2002.
22. Menke KH, Raab L, Salewski A, Steingass H, Fritz D, Schneider W. The estimation of the digestibility and metabolizable energy content of ruminant feeding stuffs from the gas production when they are incubated with rumen liquor *in vitro*. J Agric Sci Cambridge 1979;93:217-22.
23. Makkar HPS, Blümmel M, Becker K. Formation of complexes between polyvinyl pyrrolidones or polyethylene glycols and tannins and their implications in gas production and true digestibility in *in vitro* techniques. Br J Nutr 1995;73:897-933
24. Velazquez AE, Elghandour MMY, Adegbeye MJ, Pilego AB, Vallejo LH, Salem AZM, Salazar MC. Influence of dietary inclusion with corn and soybean oils, in combination with live yeast culture, on horse fecal methane, carbon dioxide and hydrogen production. J Equine Vet Sci 2019;74:42-50
25. Velázquez AE, Kholif AE, Elghandour MMY, Salem AZM, de Oca Jiménez RM, Pliego AB, Odongo N, Bórquez JL, Cipriano M, Olivares J. Effect of partial replacement of steam rolled corn with soybean hulls or prickly pear cactus in the horse's diet in the presence of live *Saccharomyces cerevisiae* on *in vitro* fecal gas production. J Equine Vet Sci 2016;42:94–101.
26. Cone JW, Van Gelder AH, Bachmann H. Influence of inoculum source dilution and storage of rumen fluid on gas production profiles. In: Gas Production: Fermentation Kinetics for Feed Evaluation and to Assess Microbial Activity. Proceedings of the EAAP Satellite Symposium on Gas Production, Wageningen, The Netherlands Proc Br Soc Anim Sci 200; pp. 15-16.

27. Faniyi TO, Adegbeye MJ, Elghandour MMY, Pilego AB, Salem AZM, Olaniyi TA, Adediran O, Adewumi MK. 2019. Role of diverse fermentative factors towards microbial community shift in ruminants. *J Appl Microbiol* 2019;30:1-10.
28. Russell WR, Gratz SW, Duncan SH, Holtrop G, Ince J, Scobbie L, Duncan G, Johnstone AM, Lobley GE, Wallace RJ, Duthie GG, Flint HJ. High-protein, reduced-carbohydrate weight-loss diets promote metabolite profiles likely to be detrimental to colonic health. *Am J Clin Nutr* 2011;93:1062-72.
29. den Besten G, van Eunen K, Groen AK, Venema K, Reijngoud DJ, Bakker BM. The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *J Lipid Res* 2013;54:2325-40.
30. Elghandour MMY, Kholif AE, Salem AZM, de Oca RM, Barbabosa A, Mariezcurrena M, Olafadehan OA. Addressing sustainable ruminal methane and carbon dioxide emissions of soybean hulls by organic acid salts. *J Clean Prod* 2016;135:194-200.
31. Monteiro ALG, da Fonseca Faro AMC, Peres MTP, Batista R, Poli CHE, Villalba JJ. The role of small ruminants on global climate change. *Acta Scient Anim Sci* 2018;40:1-11.
32. Laukova´ A, Marekova´ M. Antimicrobial spectrum of bacteriocin-like substances produced by rumen staphylococci. *Folia Microbiol* 1993;38:74–6.
33. Sung C, Kim BG, Kim S, Joo HS, Kim PI. Probiotic potential of *Staphylococcus hominis* MBBL 2–9 as anti-*Staphylococcus aureus* agent isolated from the vaginal microbiota of a healthy woman. *J Appl Microbiol* 2009;108:908–16.

34. Mauriello G, Casaburi A, Blaiotta G, Villani F. Isolation and technological properties of coagulase negative staphylococci from fermented sausages of Southern Italy. *Meat Sci* 2004;67:149–58.
35. Casaburi A, Blaiotta G, Mauriello G, Pepe O, Villani F. Technological activities of *Staphylococcus carnosus* and *Staphylococcus simulans* strains isolated from fermented sausages. *Meat Sci* 2005; 71:643–50.
36. Sakthivel PC, Kamra DN, Agarwal N, Chaudhary LC. Effect of sodium nitrate and nitrate reducing bacteria on *in vitro* methane production and fermentation with buffalo rumen liquor. *Asian-Aust J Anim Sci* 2012;25:812–7.
37. Elghandour MMY, Vázquez Chagoyán JC, Salem AZM, Kholif AE, Castañeda JSM, Camacho LM, Buendía G. *In vitro* fermentative capacity of equine fecal inocula of 9 fibrous forages in the presence of different doses of *Saccharomyces cerevisiae*. *J Equine Vet Sci* 2014;34:619–25
38. Elghandour MMY, Kholif AE, López S, Mendoza GD, Odongo NE, Salem AZM. *In vitro* gas, methane, and carbon dioxide productions of high fibrous diet incubated with fecal inocula from horses in response to the supplementation with different live yeast additives. *J Equine Vet Sci* 2016;38:64–71
39. Kholif AE, Baza-García LA, Elghandour MMY, Salem AZM, Barbabosa A, Dominguez-Vara IA, Sanchez-Torres JE. *In vitro* assessment of fecal inocula from horses fed on high-fiber diets with fibrolytic enzymes addition on gas, methane, and carbon dioxide productions as indicators of hindgut activity. *J Equine Vet Sci* 2016;39:44–50