


SEP

POLIBOTÁNICA

ISSN 1405-2768



Enero 2019

Núm. 47

POLIBOTÁNICA



Núm. 47



Enero 2019

PÁG.

CONTENIDO

- 1 Diversidad de helechos y licopodios en el área privada de conservación Talhpan, Papantla, Veracruz, México.
Diversity of ferns and lycopods in the Talhpan private conservation area, Papantla, Veracruz, Mexico.
F. Barrios-Morales | J.L. Alanís-Méndez | C.I. Carvajal-Hernández
- 13 Análisis de la estructura poblacional del frijol ayocote (*Phaseolus coccineus* L.) mediante AFLP.
Population structure analysis of ayocote beans (*Phaseolus coccineus* L.) by AFLP.
Ruiz-Salazar, R. | N. Mayek-Pérez | M.L.P. Vargas-Vázquez | S. Hernández-Delgado | J.S. Muruaga-Martínez
- 25 Diversidad, estructura arbórea e índice de valor de importancia en un bosque urbano de la Ciudad de México.
Diversity, tree structure, and importance value index in an urban park in Mexico city.
L. de L. Saavedra-Romero | P. Hernández-de la Rosa | D. Alvarado-Rosales | T. Martínez-Trinidad | J. Villa-Castillo
- 39 Heterogeneidad estructural del manglar como respuesta a factores ambientales y antrópicos en el Soconusco, Chiapas, México.
Structural heterogeneity of mangroves as a response of environmental and anthropic factors in Soconusco, Chiapas, Mexico.
E. I. Romero-Bermy | C. Tovilla-Hernández | N. Torrescano-Valle | B. Schmook
- 59 Descomposición de la madera de roble (*Quercus* spp.) in vivo e in vitro.
In vivo and in vitro wood decay of oak (*Quercus* spp.).
J. Carranza-V. | J.F. Di Stefano-G. | W. Marín-M. | M. Mata-H.
- 77 Morfología y germinación de *Chamaedorea glaucifolia* (Arecaceae), especie en peligro y endémica de México.
Morphology and germination of *Chamaedorea glaucifolia* (Arecaceae), endangered and endemic species of Mexico.
C. Orantes-García | O. Farrera-Sarmiento | R. Gutiérrez-González, | R. A. Moreno-Moreno | E. R. Garrido-Ramírez | S. J. Reyes-Zambrano
- 89 Ácido rosmarínico, fenólicos totales y capacidad antioxidante en tres variedades de *Ocimum basilicum* L. con diferentes dosis de potasio.
Rosmarinic acid, total phenolics and antioxidant capacity in three varieties of *Ocimum basilicum* L. with different potassium doses.
M.G. Ramírez-Aragón | V.J. Borroel-García | L. Salas-Pérez | J.D. López-Martínez | M.A. Gallegos-Robles | H.I. Trejo-Escareño
- 99 Efecto de extractos crudos de ajo (*Allium sativum*) sobre el desarrollo in vitro de *Aspergillus parasiticus* y *Aspergillus niger*.
Effect of garlic extracts (*Allium sativum*) on the development in vitro of *Aspergillus parasiticus* and *Aspergillus niger*.
K.G. Juárez-Segovia | E.J. Díaz-Darcía | M.D. Méndez-López | M.S. Pina-Canseco | A.D. Pérez-Santiago | M.A. Sánchez-Medina
- 113 Evaluación in vitro de la actividad citotóxica y antitumoral de plantas medicinales recomendadas en Cuetzalan del Progreso, Puebla, México.
In vitro evaluation of cytotoxic and antitumor activity of medicinal plants recommended in Cuetzalan del Progreso, Puebla, Mexico.
M.C.G. Avelino-Flores | M.D. Bibbins-Martínez | V. Vallejo-Ruiz | J. Reyes-Leyva
- 137 Partial characterization of the potentially bioactive protein fraction of *Solanum marginatum* L. f.
Caracterización parcial de una fracción proteica potencialmente bioactiva de *Solanum marginatum* L. f.
J. Guzmán-Ceferino | L.E. Cobos-Puc | C.A. Sierra-Rivera | J.C. Contreras-Esquivel | T. Durán-Mendoza | S.Y. Silva Belmares
- 153 Density and integral use evaluation of three leguminous trees in silvopastoral systems in the tropic of Guerrero, Mexico.
Evaluación de la densidad y uso integral de tres árboles leguminosos en los sistemas silvopastoriles en el trópico de Guerrero, México.
S. Rojas-Hernández | J. Olivares-Pérez | A.J. Chay-Canul | A.E. Villa-Mancera | F. Avilés-Nova | A. Córdova-Izquierdo | M.A. Damián-Valdez
- 167 Plantas para afecciones respiratorias comercializadas en tres mercados de la ciudad de Santiago de Querétaro.
Plants to treat respiratory ailments that are commercialized in three markets in Queretaro city.
J.C. Juárez-Pérez | J.A. Cabrera-Luna
- 179 Principio de conservación: coexistencia entre diversidad de especies comestibles y conocimiento tradicional.
Principle of conservation: coexistence between diversity of edible species and traditional knowledge.
X. Jasso-Arriaga



Portada

Allium sativum L. Amaryllidaceae. “Ajo”. Especie de importancia económica, extensamente cultivada y muy variable en cuanto a su morfología y fisiología. Es altamente probable que sea originaria de Asia occidental e introducida desde allí en el Mediterráneo —y luego a otras zonas del mundo— donde se cultiva desde hace más de 7 000 años.

Allium sativum L. Amaryllidaceae. “Garlic”. Species of economic importance widely cultivated and very variable in terms of its morphology and physiology. It is highly probable that it originated in western Asia and was introduced from there into the Mediterranean - and then into other parts of the world - where it has been cultivated for more than 7 000 years.

por/by **Rafael Fernández Nava**

INDEXACIÓN DE POLIBOTÁNICA

POLIBOTÁNICA se encuentra indexada en los siguientes índices internacionales:

CONACYT, índice de Revistas Mexicanas de Investigación Científica y Tecnológica del Consejo Nacional de Ciencia y Tecnología.

DOAJ, Directorio de Revistas de Acceso Público, Este servicio es de acceso gratuito a los textos completos de los artículos publicados en Revistas Científicas de excelencia académica Bibliotecas de la Universidad de Lund. Suecia.

Open Science Directory, Este directorio permite acceder a la literatura científica de aproximadamente 13000 revistas científicas. Ha sido posible gracias a programas desarrollados por numerosas universidades que permite el acceso al conocimiento a numerosos países en vías de desarrollo, desarrollado por la Biblioteca de la Universidad de Hasselt, Bélgica.

Index Copernicus International (IC), proporciona servicios bibliográficos en los que se incluyen el acceso libre a resúmenes y textos completos de los artículos científicos, así como una clasificación y el cálculo de un Factor de Descarga (DF).

CAB International

The Kew Record of Taxonomic Literature

REDALYC, Red de Revistas Científicas de América Latina y el Caribe, España y Portugal que bajo el lema "la ciencia que no se ve no existe" busca contribuir al fortalecimiento de la comunicación; científica de Iberoamérica

LATINDEX, Sistema regional de información en línea para revistas científicas de América Latina, el Caribe, España y Portugal. Latindex Es producto de la cooperación de una red de instituciones que funcionan de manera coordinada para reunir y diseminar información bibliográfica sobre las publicaciones científicas seriadas producidas en la región.

PERIODICA, (Índice de Revistas Latinoamericanas en Ciencias).

SciELO, está conformada por las colecciones de revistas académicas de 15 países: Argentina, Bolivia, Brasil, Chile, Colombia, Costa Rica, Cuba, España, México, Paraguay, Perú, Portugal, Sudáfrica, Uruguay y Venezuela.

Dialnet es un portal de difusión de la producción científica hispana. Su base de datos, de acceso libre, fue creada por la Universidad de La Rioja (España) y constituye una hemeroteca virtual que contiene los índices de las revistas científicas y humanísticas de España, Portugal y Latinoamérica. El texto completo de muchos de los documentos está disponible en línea. En el portal colaboran numerosas universidades españolas e hispanoamericanas que realizan los volcados de sumarios de revistas. También incorpora bases de datos con documentos en otros idiomas.

REDIB Red Iberoamericana de innovación y conocimiento científico.



INSTITUTO POLITÉCNICO NACIONAL

Director General: *Dr. Mario Alberto Rodríguez Casas*

Secretario General: *Dr. Héctor Leoncio Martínez Castuera*

Secretario Académico: *Dr. Jorge Toro González*

Secretario de Extensión e Integración Social: *Dr. Luis Alfonso Villa Vargas*

Secretario de Investigación y Posgrado: *Dr. Juan Silvestre Aranda Barradas*

Secretaria de Servicios Educativos: *Mtra. María Guadalupe Vargas Jacobo*

Secretario de Administración: *C.P. Jorge Quintana Reyna*

Directora de Educación Superior: *M. en C. Rosalía María del Consuelo Torres Bezaury*

ESCUELA NACIONAL DE CIENCIAS BIOLÓGICAS

Director:

Dr. Gonzalo Trujillo Chávez

Subdirectora Académica:

M. en C. Martha Patricia Cervantes Cervantes

Subdirector de Servicios Educativos e Integración Social:

Dr. Omar Hernández Montes

Subdirector Administrativo: *Ing. Raúl Chávez Alvircio*

Jefe de la Sección de Estudios de Posgrado e Investigación:

Dr. Gerardo Aparicio Ozores

Jefe del Departamento de Extensión de la Cultura:

Ing. Amado Enrique Takeshi Saiki Shirai

POLIBOTÁNICA, Año 21, No. 47, enero-junio 2019, es una publicación semestral editada por el Instituto Politécnico Nacional, a través de la Escuela Nacional de Ciencias Biológicas. Unidad Profesional Lázaro Cárdenas, Prolongación de Carpio y Plan de Ayala s/n, Col. Santo Tomas C.P. 11340 Delegación Miguel Hidalgo México, D.F. Teléfono 57296000 ext. 62331. <http://www.herbario.encb.ipn.mx/>, Editor responsable: Rafael Fernández Nava. Reserva de Derechos al Uso Exclusivo del Título No. 04-2015-011309001300-203. ISSN impreso: 1405-2768, ISSN digital: 2395-9525, ambos otorgados por el Instituto Nacional del Derecho de Autor. Responsable de la última actualización de este número, Unidad de informática de la ENCB del IPN, Rafael Fernández Nava, Unidad Profesional Lázaro Cárdenas, Prolongación de Carpio y Plan de Ayala s/n, Col. Santo Tomas C.P. 11340 Delegación Miguel Hidalgo México, D.F.

Las opiniones expresadas por los autores no necesariamente reflejan la postura del editor de la publicación.

Queda estrictamente prohibida la reproducción total o parcial de los contenidos e imágenes de la publicación sin previa autorización del Instituto Politécnico Nacional.

REVISTA BOTÁNICA INTERNACIONAL DEL INSTITUTO POLITÉCNICO NACIONAL

EDITOR EN JEFE

Rafael Fernández Nava

EDITORA ASOCIADA

María de la Luz Arreguín Sánchez

COMITÉ EDITORIAL INTERNACIONAL

Christiane Anderson
University of Michigan
Ann Arbor, Michigan, US

Heike Vibrans
Colegio de Postgraduados
Estado de México, México

Graciela Calderón de Rzedowski
Instituto de Ecología del Bajío
Pátzcuaro, Mich., México

Theodore S. Cochran
University of Wisconsin
Madison, Wisconsin, US

Hugo Cota Sánchez
University of Saskatchewan
Saskatoon, Saskatchewan, Canada

Fernando Chiang Cabrera
Universidad Nacional Autónoma de México
Ciudad de México, México

Thomas F. Daniel
California Academy of Sciences
San Francisco, California, US

Francisco de Asis Dos Santos
Universidad Estadual de Feira de Santana
Feira de Santana, Brasil

Carlos Fabián Vargas Mendoza
Instituto Politécnico Nacional
Ciudad de México, México

José Luis Godínez Ortega
Universidad Nacional Autónoma de México
Ciudad de México, México

José Manuel Rico Ordaz
Universidad de Oviedo
Oviedo, España

Edith V. Gómez Sosa
Instituto de Botánica Darwinion
Buenos Aires, Argentina

Jorge Llorente Bousquets
Universidad Nacional Autónoma de México
Ciudad de México, México

Delia Fernández González
Universidad de León
León, España

Jerzy Rzedowski Rotter
Instituto de Ecología del Bajío
Pátzcuaro, Mich., México

Luis Gerardo Zepeda Vallejo
Instituto Politécnico Nacional
Ciudad de México, México

Claude Sastre
Muséum National d'Histoire Naturelle
Paris, Francia

Mauricio Velayos Rodríguez
Real Jardín Botánico
Madrid, España

Noemí Waksman de Torres
Universidad Autónoma de Nuevo León
Monterrey, NL, México

Julieta Carranza Velázquez
Universidad de Costa Rica
San Pedro, Costa Rica

Tom Wendt
University of Texas
Austin, Texas, US

CORRECCIÓN DE ESTILO

Margarita Sam Rodríguez

DISEÑO Y FORMACIÓN ELECTRÓNICA

Luz Elena Tejeda Hernández

OPEN JOURNAL SYSTEM Y TECNOLOGÍAS DE LA INFORMACIÓN

Pedro Aráoz Palomino

Toda correspondencia relacionada con canje, compra o suscripción deberá ser dirigida a:

Dr. Rafael Fernández Nava
Editor en Jefe de

POLIBOTÁNICA

Departamento de Botánica
Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional
Apdo. Postal 17-564, CP 11410, Ciudad de México

<http://www.herbario.encb.ipn.mx/polibotanica.htm>

Correo electrónico:
rfernand@ipn.mx
polibotanica@gmail.com

POLIBOTÁNICA es una revista indexada en el Sistema de Clasificación de
Revistas Mexicanas de Ciencia y Tecnología (CRMcyT)
CONACYT

y en los índices bibliográficos internacionales
LATINDEX y Red AlyC

Suscripción anual
México \$250.00 MN
Extranjero \$35.00 USD



Polibotánica

ISSN electrónico: 2395-9525

polibotanica@gmail.com

Instituto Politécnico Nacional

México

<http://www.polibotanica.mx>

DENSITY AND INTEGRAL USE EVALUATION OF THREE LEGUMINOUS TREES IN SILVOPASTORAL SYSTEMS IN THE TROPIC OF GUERRERO, MEXICO

EVALUACIÓN DE LA DENSIDAD Y USO INTEGRAL DE TRES ÁRBOLES LEGUMINOSOS EN LOS SISTEMAS SILVOPASTORILES EN EL TRÓPICO DE GUERRERO, MÉXICO

Rojas-Hernández, S.; J. Olivares-Pérez, A.J. Chay-Canul, A.E. Villa-Mancera, F. Avilés-Nova, A. Córdova-Izquierdo, y M.A. Damián-Valdez.

DENSITY AND INTEGRAL USE EVALUATION OF THREE LEGUMINOUS TREES IN SILVOPASTORAL SYSTEMS IN THE TROPIC OF GUERRERO, MEXICO.

EVALUACIÓN DE LA DENSIDAD Y USO INTEGRAL DE TRES ÁRBOLES LEGUMINOSOS EN LOS SISTEMAS SILVOPASTORILES EN EL TRÓPICO DE GUERRERO, MÉXICO.

**DENSITY AND INTEGRAL USE EVALUATION OF THREE LEGUMINOUS TREES
IN SILVOPASTORAL SYSTEMS IN THE TROPIC OF GUERRERO, MEXICO**

**EVALUACIÓN DE LA DENSIDAD Y USO INTEGRAL DE TRES ÁRBOLES LEGUMINOSOS
EN LOS SISTEMAS SILVOPASTORILES EN EL TRÓPICO DE GUERRERO, MÉXICO**

Rojas-Hernández, S.;
J. Olivares-Pérez,
A.J. Chay-Canul,
A.E. Villa-Mancera,
F. Avilés-Nova,
A. Córdova-Izquierdo,
y M.A. Damián-Valdez

DENSITY AND INTEGRAL
USE EVALUATION OF
THREE LEGUMINOUS
TREES IN SILVOPASTORAL
SYSTEMS IN THE TROPIC
OF GUERRERO, MEXICO

EVALUACIÓN DE LA
DENSIDAD Y USO INTEGRAL
DE TRES ÁRBOLES
LEGUMINOSOS EN LOS
SISTEMAS
SILVOPASTORILES EN EL
TRÓPICO DE GUERRERO,
MÉXICO

POLIBOTÁNICA

Instituto Politécnico Nacional

Núm. 47: 153-166. Enero 2019

DOI:

10.18387/polibotanica.47.11

S. Rojas-Hernández

J. Olivares-Pérez/ olivaares@hotmail.com

*Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma de Guerrero,
Ciudad Altamirano, Guerrero, México.*

A.J. Chay-Canul

*Facultad de Medicina Veterinaria, Universidad Juárez Autónoma de Tabasco,
Villahermosa, Tabasco, Mexico*

A.E. Villa-Mancera

*Facultad de Medicina Veterinaria y Zootecnia,
Benemérita Universidad Autónoma de Puebla, Tecamachalco Puebla, México*

F. Avilés-Nova

*Universidad Autónoma del Estado de México, Toluca,
Estado de México-Centro 8 Universitario, Temascaltepec, México*

A. Córdova-Izquierdo

Facultad de Medicina Veterinaria y Zootecnia-UAM-X, México

M.A. Damián-Valdez

*Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma de Guerrero,
Ciudad Altamirano, Guerrero, México.*

RESUMEN: En el Trópico Mexicano se estudió la distribución, densidad y usos de los árboles *Pithecellobium dulce* Roxb Benth, *Gliricidia sepium* (Jacq) Steud, *Haematoxylum brasiletto* Karst y se agregó su follaje a las dietas de cabritos y midió la respuesta productiva y la digestibilidad aparente. Los ganaderos dieron seis usos (leña, postes, sombra, medicinales, consumo humano y artesanal). Los árboles estuvieron en cercas vivas y dispersos en potreros con densidades de 4,87 árboles 100⁻¹ metros lineales y 1,79 árboles ha⁻¹; *P. dulce* Roxb Benth tuvo mayor talla y se localizó entre 250 a 1332 msnm. Las cenizas (P<0,0001) y fibras detergentes (P<0,01) del *P. dulce* (T1) fueron más digestibles. La CA y GDP entre los cabritos fueron diferentes entre tratamientos. La temperatura rectal sólo se afectó (P<0,0001) por la hora de evaluación. El pH ruminal de los cabritos fue diferente entre tiempos de evaluación (P<0,0001) y tratamientos (P<0,0003). Se concluye que la densidad de árboles es baja y podría tener un impacto sobre la fertilidad del suelo y la aportación de biomasa para la alimentación animal; La digestibilidad de las dietas y respuesta productiva fueron más eficientes en los animales alimentados con *P. dulce* Roxb Benth (T1).

Palabras clave: alimentación, leguminosas, pH, rumiante, digestibilidad aparente

ABSTRACT: In the Tropic of Guerrero Mexico the distribution, the density and uses of trees *Pithecellobium dulce* Roxb Benth, *Gliricidia sepium* (Jacq) Steud,

Haematoxylum brasiletto Karst were studied and its foliage was added to the diets of kids and productive response and apparent digestibility was measured. The livestock farmers have six complementary uses (firewood, poles, shade, medicinal, human consumption and artisanal). The species were identified in live fences and scattered in paddocks with densities of 4.87 trees per 100 linear meters and 1.79 trees ha⁻¹; *P. dulce* was the largest size and identified from 250 to 1332 masl. The ashes (P<0.0001) and detergent fibre (P<0.01) of *P. dulce* (T1) had higher apparent digestibility. The feed conversion and daily weight gain of the kids were different due to the effect of foliage trees. The rectal temperature was only affected (P<0.0001) by the time of evaluation. The ruminal pH of kids was affected by time evaluation (P<0.0001) and the treatments (P<0.0003). It is concluded that the density of trees is low and could be of impact on soil fertility and the contribution of biomass for animal feed; the apparent digestibility of diets and productive response were more efficient in the animals fed with *P. dulce* (T1).

Key words: feeding, legumes, pH, ruminant, apparent digestibility.

INTRODUCTION

Fodder trees in silvopastoral systems are an important component from the environmental point of view that can reduce soil erosion, to act as a carbon sink and integrate great amounts of organic matter and nitrogen to the soil. Moreover backgrounds suggest that in cattle ranches the trees can be used for various services such as natural shade, for obtaining wood (beams, wooden posts and poles for rural construction), as medicine and food for animals (Olivares-Pérez *et al.*, 2016). Has been shown that the foliage can be used in the control of intestinal parasites and thus increase the productivity in goats (León-Castro *et al.*, 2015) this because tannins are compounds found in the leaves and / or fruit of the trees and they can have an effect against parasites (Patra & Saxena, 2011; Rojas-Hernández *et al.*, 2016; Williams *et al.*, 2014). The density of the fodder trees is important in the prairies, if the density is high can contribute to maintain and / or restore degraded soils (Olivares-Pérez *et al.*, 2016; Reis *et al.*, 2010), in addition, the contribution of biomass useful as animal feed is higher as well (Rojas-Hernández *et al.*, 2016). The biomass of trees as an alternative feed for ruminants has been used in several studies with positive results in animal response (the dry matter intake, weight gain, feed conversion and diet digestibility) (Olivares *et al.*, 2013; Olivares Perez *et al.*, 2013; Rojas-Hernandez *et al.*, 2015; Rojas Hernández *et al.*, 2015). However a constraint to use the arboreal resource as food is its tannin content, these compounds in high concentrations can confer organoleptic characteristics to food that reduce the palatability and consumption of dry matter in animals (Lee *et al.*, 2015; Patra & Saxena, 2011). In addition, reports indicate that tannins have defaunate effect and are capable of binding to dietary nutrients (protein, fiber and carbohydrates) and affect digestibility and reduce animal productivity (Lorenz *et al.*, 2014; Patra & Saxena, 2011). It has also been reported that changes in ruminal pH outside its range may have an impact on the health of ruminant animals and the factor most closely related is the food type they receive in the diet (Chibisa *et al.*, 2016; Lee *et al.*, 2015; León-Castro *et al.*, 2015). The objectives were to study the distribution, density and uses of trees *Pithecellobium dulce* Roxb Benth, *Gliricidia sepium* Jacq Steud and *Haematoxylum brasiletto* Karst and add foliage to diets for kids and measure the productive response and apparent digestibility.

MATERIAL AND METHODS

Location of the experimental site

The study was conducted in the Tropical region of Guerrero, Mexico located at 18° 20' 30" NL and 100° 39' 18" WL. The climate that prevails is the Aw0, is the driest of the sub humid warm with summer rains. The average temperature and rainfall are of 28°C and 1010.7 mm.

Descriptive study

Location, distribution, uses and dasometric measures: The information on uses of trees was obtained by direct interviews 81 livestock farmers in the study area. The location and distribution was assessed and measures dasometric trees were recorded in four plots transects at each ranch. Each transect was one ha for scattered trees in pastures and hundred linear meters (L) for trees in live fences.

Experimental study

Foliage Collection: The foliage of *Pithecellobium dulce* Roxb Benth, *Haematoxylum brasiletto* Jacq Steud and *Gliricidia sepium* Karst were collected manually by pruning trees and dehydrated by exposure to solar rays to constant weight (on average 88% of dry matter). The dehydrated foliage was milled in a hammer mill to a particle size of five mm on average and were integrated to rations of goat kids.

Management of animals

Experimental animals: 15 weaned male Creole kids were used with an average, average live weight of 14 ± 3 kg from existing herds in the study area, used to graze the native vegetation from birth. The animals were kept in individual pens with a vital space (2 m^2), with free access to drinking trough and manger where treatments were given. At the beginning of the experiment the kids were dewormed subcutaneously with ivermectin at doses of 0.2 mg / kg body weight (Sumano & Ocampo, 2006).

Treatment assignment: Kids are distributed through a completely randomized design in three treatments that were provided ad libitum for 68 days (8 days of adaptation and 60 days of evaluation). Treatments were, isoproteic (11.0%) and isocaloric diets (10.5 Mj ME kg DM) added with 30% of foliage *Pithecellobium dulce* Roxb Benth (T1), *Gliricidia sepium* Jacq Steud (T2) and *Haematoxylum brasiletto* Karst (T3) (table 1) prepared according to the nutritional requirements of animals (Nutritional Research Council (NRC), 2007). The final diets were analyzed for content of acid (ADF) and neutral (NDF) detergent fiber with the Van Soest and Wayne method (Van Soest & Lewis, 1991) and the ash content (Ce), organic matter (OM) and crude protein (CP) according to the AOAC methods (Association of Official Analysis Chemists International (AOAC), 2000).

Measures variables

Animal response: Nutrient intake (dry matter, organic matter, protein, neutral detergent fiber and acid) were measured and obtained by multiplying the daily consumption of DM by the content of each nutrient in the diet (table 1). Weight change and dry matter intake were recorded. The live weight was measured with an electronic scale with capacity of 500 kg and graduated in grams; the dry matter intake (DMI) was determined by subtracting the amount offered from the amount rejected daily. The daily weight gain (DWG) was estimated by subtracting the final weight, the initial weight and divided between the days of the study. The feed conversion (FC) was calculated by dividing the feed intake between the DWG.

Rectal temperature: The inclusion of foliage of trees in diets that constituted the treatments, can bring toxic substances for goats (condensed tannins, CT) that in some situations compromise the thermoregulatory capacity of the animals. For this reason, during the last seven days of the experiment with the animals of each treatment the rectal temperature was measured at three times (07:00, 13:00 and 18:00 h) with an electronic thermometer graduated in degrees Celsius.

Apparent digestibility of nutrients from the diet: For seven days, all animals' faeces were collected using harnesses collectors placed on each adapted animal during fourteen days. For the collected faeces the content of DM, OM, CP, ADF and NDF were determined. Known nutrient content in the diet and faeces and the apparent digestibility of the diet (DM, OM, CP, ADF and NDF) were calculated, using the equation (International Livestock Centre for Africa (ILCA), 1990):

$$\text{Digestibility (\%)} = \frac{\text{Consumption} - \text{Fecal excretion}}{\text{Consumption}} \times 100$$

Where, Digestibility (%) is the nutrient digestibility, consumption is equal to consumed nutrient, faecal excretion represents each nutrient excreted in the faeces.

Measuring of ruminal hydrogen potential (pH): On the last day of the experiment of each animal 60 mL of ruminal fluid were collected at different times (0, 1, 3 and 6 h post feeding), immediately after collection using a potentiometer was measured ruminal pH fluctuation as indicative of the environmental condition for the fermentative rumen microflora activity.

Table 1. Ingredients and chemical composition of treatments with fruits, dry basis quantities (DM)

Ingredient (kg / 100 kg DM)	T1	T2	T3	SEM
Foliage of <i>P. dulce</i>	30.0			
Foliage of <i>G. sepium</i>	-	30.0	-	-
Foliage of <i>H. brasiletto</i>	-	-	30.0	-
Ground corn	6.0	0.0	-	-
Ground cob	61.0	65.0	61.0	-
Soybean meal	2.0	4.0	8.0	-
Mineral premix	1.0	1.0	1.0	-
Totals DM (kg)	100.0	100.0	100.0	-
	Chemical composition (% DM)			
Crude protein	11.7	12.0	12.6	-
Ashes	3.9	3.9	2.9	2.6
Organic matter	96.0	96.0	97.0	2.7
Neutral detergent fibre	33.3	27.5	38.3	5.6
Acid detergent fibre	14.6	13.7	17.8	0.5
ME Mj / kg DM	11.3	11.0	11.4	-
<i>In vitro</i> gas production (96 h, mL / g DM)	277.8 ^a	236.0 ^b	235.4 ^b	5.8

T1: *P. dulce* 30%; T2: *G. sepium* 30%; T3: *H. brasiletto* 30% SEM: standard error of the means

Statistical analysis

Descriptive study: The location, distribution, uses, density and trees dasometric measures by descriptive statistics (Steel & Torrie, 2006).

Experimental study: The data of variables of animal response (dry matter intake and nutrient apparent digestibility, daily weight gains and feed conversion) were analysed by variance according to a completely randomized design with the statistical model $Y_{ij} = \mu + \tau_i + \zeta_{ij}$, where Y_{ij} is the response variable, μ is the overall mean, τ_i treatment effect i (1, 2 and 3 treatments) and ζ_{ij} is random error attributed to the i th treatment in the j th repetition (1, 2 ... 5 male kids).

The rectal temperature and ruminal pH data were analysed in a completely randomized design in 3 x 3 factorial arrangement (three forages for three times as rectal temperature, 7:00, 13:00 and 18:00 respectively) and 3 x 4 (three times foliage's four ruminal pH measurement, 0, 1, 3 and 6 h post-feeding, respectively); statistical model $Y_{ijk} = \mu + A_i + B_j + (AB)_{ij} + \zeta_{ijk}$; Where: Y_{ijk} = rectal temperature, μ is the general mean; A_i is the treatment effect ($i = 1, 2$ and 3 treatments) is the measured times B_j effect; $A * B_{ij}$ is the first-order interaction and ζ_{ijk} is the mistake in terms of n-1 ($\sigma^2, 0$) (SAS, 2002).

Correlation analysis (r) were developed to determine the relationship between measurement times (hours) with ambient temperature (AT) and rectal temperature (RT) of animals and regression analysis (r^2) were performed between times measurement and AT with RT animal. The averages of the evaluated variables were compared with the Tukey test ($P < 0.05$) (SAS, 2002).

RESULTS

Descriptive study

Location, density and uses of trees: In figure 1, letter "a" shows that the *P. dulce* Roxb Benth adapts to a wide range of altitudes from 250 masl to 1 332 masl compared with the trees of *G. sepium* Jacq Steud and *H. brasiletto* Karst that were located up to 455 and 512 masl respectively. In addition, the three trees are distributed in arrangement of live fences and scattered trees in 85.3 and 75.5% of the transects evaluated in livestock ranches (fig. 1, item b), with an average density of 4.87 ± 1.4 trees per 100 m linear and in live fences of 1.79 ± 0.5 trees per ha in the rangeland (fig. 1, item c).




Fig. 1. Location (masl), distribution and density of forage trees in livestock ranches in the Tropical Region of Guerrero Mexico.

In addition to the feed, identified six complementary uses were (firewood, poles, shade, medicinal, human consumption and artisanal) for which farmers use *P. dulce* Roxb Benth, four at *H. brasiletto* Karst (firewood, poles, shade, and medicinal) and only two for *G. sepium* Jacq Steud (shade and medicinal). Although the *P. dulce* Roxb Benth has a greater dasometric size, the trees of *H. brasiletto* Karst and *G. sepium* Jacq Steud reach a development that allows them to be used equally as shade for cattle in pens and grazing areas (table 2). The medicinal uses were to counter diabetes (*G. sepium* Jacq Steud) for the kidney (*H. brasiletto* Karst) and as dewormer (*P. dulce* Roxb Benth) (fig. 2).

Table 2. Dasometric measures and complementary uses of fodder trees in livestock ranches in the Tropical Region of Guerrero, Mexico.

Scientific name	<i>P. dulce</i>	<i>G. sepium</i>	<i>H. brasiletto</i>
Dasometric measures			
Basal diameter (cm)	128	76.4	62.0
Diameter breast height (cm)	95.7	62.1	58.8
Height (m)	12.3	9.2	8.3
Additional uses of trees			
Firewood	✓	-	✓
Posts	✓	-	✓
Shade	✓	✓	✓
Medicinal	✓	✓	✓
Human consumption	✓	-	-
Handicraft	✓	-	-




Fig. 2. Main medicinal uses of fodder trees in livestock ranches in the Tropical Region of Guerrero, Mexico.

Experimental study

Animal response: The nutrient intake (DM, NDF, ADF and CP) was not different ($P > 0.05$) among kids who received the different treatments (T1, T2 and T3) (Table 3). The daily weight gain was higher ($P < 0.002$) in kids fed with the T1 (127.4 g) and T2 (89.0 g) added with foliage *P. dulce* Roxb Benth and *G. sepium* Jacq Steud, respectively, and feed conversion was higher (4.8 kg DM) ($P < 0.03$) in the kids fed with the diet added with foliage of *P. dulce* Roxb Benth (T1).

The values of apparent digestibility of the different diets (T1, T2 and T3), show that the NDF and the ashes contained in the diets were more utilized by kids of T1 with foliage of *P. dulce* Roxb Benth, additionally this relates with the biggest weight gain and best feed conversion shown by animals (table 3).

Table 3. Consumption of dry matter and nutrients, daily weight gain and feed conversion of growth kids fed diets containing foliage of *P. dulce*, *G. sepium* and *H. brasiletto*.

Treatments	T1	T2	T3	SEM	Significance
Dry matter intake (g / day)	606.5	567.6	594.3	57.5	0.71
Crude protein intake (g / day)	70.9	68.1	74.8	22.6	0.96
Neutral detergent fibre intake (g / day)	201.9	156.1	227.6	62.3	0.41
Acid detergent fibre intake (g / day)	88.5	77.7	105.8	28.4	0.51
Metabolizable energy intake (Mj / day)	6.8	6.2	6.7	1.8	0.20
Daily weight gain (g)	124.7 ^a	89.0 ^a	47.0 ^b	15.5	0.002
Feed conversion (g)	4.8 ^a	6.3 ^{ab}	13.9 ^b	3.5	0.03
Apparent digestibility (%)					
Dry matter	73.5	73.6	72.6	2.9	0.90
Organic matter	75.3	74.0	73.1	2.9	0.66
Acid detergent fibre	38.0 ^a	25.4 ^b	51.5 ^a	7.4	0.01
Neutral detergent fibre	57.3 ^a	39.6 ^b	54.2 ^a	5.2	0.01
Crude protein	73.1	73.6	72.3	3.5	0.90
Ashes	61.4 ^a	57.6 ^a	28.7 ^b	3.5	0.0001

T1: *P. dulce* 30%; T2: *G. sepium* 30%; T3: *H. brasiletto* 30% SEM: Standard error of the means; ^{abc} Different literal values in the same row are different, Tukey ($P < 0.05$).

Rectal temperature: The rectal temperature of the animals did not differ by treatment ($P > 0.210$), however an effect of evaluation time ($P < 0.0001$) (table 4) was observed. The rectal temperature gradually increased during daylight hours and ranged between 38.4 ° and 38.7°C at 07:00 h to 39.3° and 39.5°C for 18 h. Figure 3 shows that the day brightness hours is correlated (r) with environmental temperature and the rectal temperature of the animals; furthermore the regression analysis (r^2) shows that the variation in rectal temperature of the kids depend directly on the assessment time and environmental temperature. *Ruminal pH:* The ruminal pH was higher ($P < 0.0003$) in kids fed with the T1 with 6.49 on average in relation to animals of the T2 and T3. It was also observed that the pH gradually decreased ($P < 0.0001$) six hours after feeding the animals, ie the pH oscillate between 6.51 and 7.07 at zero hour and between 6.24 and 6.42 to the sixth hour after feeding (table 4). In addition to the independent effects, interactive effect were observed ($P < 0.003$) between treatments by time (hours) pH assessment after feeding the animals.

Table 4. Rectal temperature and ruminal pH behaviour of kids fed diets added with fodder trees foliage

Treatment diet	Time (hours)	Rectal temperature (°C)	*Time (hours)	Ruminal pH
T1 (30% foliage of <i>P. dulce</i>)	07:00	38.7 ^{bc}	0	7.07 ^a
	13:00	39.1 ^{abc}	1	6.35 ^{bcd}
	18:00	39.5 ^a	3	6.29 ^{bcd}
			6	6.59 ^{cb}
Means		-		6.57 ⁱ
T2 (30% foliage of <i>G. sepium</i>)	07:00	38.7 ^{bc}	0	6.63 ^b
	13:00	38.9 ^{ab}	1	6.24 ^{cd}
	18:00	39.3 ^{ab}	3	6.23 ^d
			6	6.28 ^{bcd}
Means		-		6.34 ⁱⁱ
T3 (30% foliage of <i>H. brasiletto</i>)	07:00	38.4 ^c	0	6.51 ^{bcd}
	13:00	38.9 ^{ab}	1	6.32 ^{bcd}
	18:00	39.3 ^{ab}	3	6.39 ^{bcd}
			6	6.42 ^{bcd}
Means		-		6.41 ⁱⁱ
Standard error of the means		0.27		0.12
Significance				
Treatments		0.210		0.0003
Time		< 0.0001		< 0.0001
Treatments * Time		0.8800		0.003

^{abcd} Values with different letters in the same column are different, Tukey (P<0.05)

^{i,ii} Differences between treatment means, Tukey (P<0.0003)

* Measurements according to the feeding schedule




Fig. 3. Correlation and regression between the independent variable time (three hours a day) and ambient temperature with a rectal temperature goat kids as the dependent variable.

DISCUSSION

Descriptive study

Location, density and uses of trees: The information in the figure 1 indicates that the presence of these trees in cattle ranches is very common, however the same density is very low compared to densities of 50 trees ha⁻¹ reported in other studies (Cajas-Giron & Sinclair, 2001). This assumes that the negative effects of the trees on pastures require control by cutting or burning, and are stronger than the beneficial effects that farmers get, however, diversified use can promote an increase in tree density (Olivares-Pérez *et al.*, 2016).

Based on the everyday complementary uses that livestock farmers give to the trees demonstrate the need to implement an integral handling of the arboreal stratum that mitigates environmental degradation, promote diversified use and increased density per surface unit, because it may represent a savings in maintenance costs of fences, construction of artificial shade, herbal medicine and in buying feed for animals among other things (Guerreiro *et al.*, 2015; Olivares-Pérez *et al.*, 2016; Tálamo *et al.*, 2015).

Experimental study

Animal response: The dry matter intake it was not different (table 3), however Olivares *et al.*, (2013) report differences in kids fed with foliage of different trees and different level of inclusion in the diet. Contrary reported no differences in dry matter intake by goats fed diets with different levels of inclusion of *Erythrina brucei* Schweinf (Yinnesu & Nurfeta, 2012).

The behaviour in FC of the animals is attributed to the greater weight gain of goat kids of T1, without requiring a significant increase in the consumption of dry matter compared to kids of the T2 and T3 (table 3). Also studies report weight gains and feed conversion higher in kids fed with foliage of *P. dulce* Roxb Benth and *G. sepium* Jacq Steud (Olivares *et al.*, 2013). In research developed with the use of biomass of trees similar results were reported in weight gain and feed conversion in ruminants (Oni *et al.*, 2008; Sanon *et al.*, 2008; Tamir & Asefa, 2009; Thang *et al.*, 2009).

The greater apparent digestibility of the ash fractions and NDF in the diet of T1 with foliage of *P. dulce* Roxb Benth (table 3) favored the weight gain and the feed conversion of the animals. The fraction of ashes due to the contribution of minerals and the fraction of NDF by the contribution of energy to the activity of microbial flora in the rumen (Hamed & Elimam, 2009; Kumara *et al.*, 2009; Olivares-Pérez *et al.*, 2013). Furthermore, it has been observed that a mineral deficiency in ruminant diet can cause a decrease in the rumen microbial growth and affect the fermentative activity of bacteria on the fibre (Feng *et al.*, 2013). Also in ruminants decreased dry matter intake and the digestibility of diets deficient in phosphorus (Feng *et al.*, 2013).

Rectal temperature: The results in rectal temperature of the animals (Table 4) indicated that the variation in this physiological constant was attributed to an increase in brightness and accumulation of heat during the day and not the caloric increase arising from the digestion of diets added with some of the foliage of the trees. The correlation and regression analyses of the Figure 3 confirm that the rectal temperature was directly related to the ambient temperature and the time of day in which the physiological constant was evaluated. Also Catanese *et al.*, (2014) who found that lambs supplemented with condensed tannins at concentrations of between 3.0 and 7.7 g / 100 g DM, did not change the rectal temperature of the animals out of their normal range 38.3 to 39.9°C (Fielder, 2015), and equally fluctuations in rectal temperature were attributed to environmental factors.

Ruminal pH: Although the results indicate differences in the ruminal pH between kids by the effect of treatment (table 4) is observed that variations, do not exceed the ranges published by different researchers in goats under different feeding regimes (Cantalapiedra-Hijar *et al.*, 2009; Paengkoum & Paengkoum, 2010; Wang *et al.*, 2009). The effects of treatment, time and interaction treatment by time on ruminal pH values kids observed, indicate that foliage of trees of *P. dulce* Roxb Benth, *G. sepium* Jacq Steud and *H. brasiletto* Karst can include up to 30% of the diet without problems or digestive imbalances. This is because the ruminal pH is an indicator of the balance of the ruminal flora, fermentative processes and end products of digestion of food (Abarghuei *et al.*, 2011; Feng *et al.*, 2013; Ondiek *et al.*, 2013; Paengkoum, 2010; Pérez-Sato *et al.*, 2011; Supreena & Paengkoum, 2011).

CONCLUSIONS

Of the study concludes that the density of three trees was low. However, although farmers the used for various services, do not have the culture for projecting an increase in the density of species assessed in live fences systems and rangeland scattered trees. It is concluded that the addition of *P. dulce* Roxb Benth foliage favoured the productive response of the goat kids in weight gain and feed conversion, greater fibre digestibility and ashes as mineralized fraction. The pH was different between animals receiving the foliage of trees *Pithecellobium dulce* Roxb Benth, *Gliricidia sepium* Jacq Steud and *Haematoxylum brasiletto* Karst in the diet, however the observed values never fluctuated outside the ranges established for small ruminants, therefore no digestive disorders were observed affecting eating behaviour, productive performance and health of the animals that destabilized thermoregulatory capacity. The body temperature was different among goat kids because the evaluation hours by the accumulation of environmental temperature caused by solar radiation during the day.

LITERATURA CITADA

- Abarghuei, M. J., Rouzbehan, Y., & Alipour, D. (2011). Effect of oak (*Quercus libani* Oliv.) leave tannin on ruminal fermentation of sheep. *Journal of Agricultural Science and Technology*, 13(SUPPL.), 1021–1032.
- Association of Official Analysis Chemists International (AOAC). (2000). *Official Methods of Analysis of AOAC International*. Association of Official Analysis Chemists International. <https://doi.org/10.3109/15563657608988149>
- Cajas-Giron, Y. S., & Sinclair, F. L. (2001). Characterization of multistrata silvopastoral systems on seasonally dry pastures in the Caribbean Region of Colombia. In *Agroforestry Systems* (Vol. 53, pp. 215–225). <https://doi.org/10.1023/A:1013384706085>
- Cantalapiedra-Hijar, G., Yáñez-Ruiz, D. R., Martín-García, A. I., & Molina-Alcaide, E. (2009). Effects of forage:concentrate ratio and forage type on apparent digestibility, ruminal fermentation, and microbial growth in goats. *Journal of Animal Science*, 87(2), 622–31. <https://doi.org/10.2527/jas.2008-1142>
- Catanese, F., Distel, R. A., & Villalba, J. J. (2014). Effects of supplementing endophyte-infected tall fescue with sainfoin and polyethylene glycol on the physiology and ingestive behavior of sheep. *Journal of Animal Science*, 92(2), 744–757. <https://doi.org/10.2527/jas.2013-6713>
- Chibisa, G. E., Beauchemin, K. A., & Penner, G. B. (2016). Relative contribution of ruminal buffering systems to pH regulation in feedlot cattle fed either low- or high-forage diets. *Animal*, 10(7), 1164–1172. <https://doi.org/10.1017/S1751731115002888>
- Feng, Y., Cong, Y., Deng, H., Guo, D., & Liu, H. (2013). Effects of 2-hydroxy-4(methylthio) butanoic acid isopropyl ester on rumen fermentation in cashmere goats. *Revista Brasileira de Zootecnia*, 42(5), 342–346. <https://doi.org/10.1590/S1516-35982013000500006>

- Fielder, S. E. (2015). *The Merck Veterinary Manual*. USA: OCEANO. Retrieved from http://www.merckvetmanual.com/mvm/appendixes/reference_guides/normal_rectal_temperature_ranges.html
- Guerreiro, M. F., Nicodemo, M. L. F., & Porfirio-da-Silva, V. (2015). Vulnerability of ten eucalyptus varieties to predation by cattle in a silvopastoral system. *Agroforestry Systems*, 89(4), 743–749. <https://doi.org/10.1007/s10457-015-9797-7>
- Hamed, A. H. M., & Elimam, E. M. (2009). Effects of Rabaa ash alkali treatment of sesame straw on chemical composition and degradation in the rumen of Nubian goats. *Pakistan Journal of Nutrition*, 8(9), 1344–1348. <https://doi.org/10.3923/pjn.2009.1344.1348>
- International Livestock Centre for Africa (ILCA). (1990). *Livestock systems research manual* (Vol. 1). Ethiopia: International Livestock Centre for Africa, Addis Ababa. Retrieved from <http://www.fao.org/Wairdocs/ILRI/x5469E/x5469e00.htm#Contents>
- Kumara Mahipala, M. B. P., Krebs, G. L., McCafferty, P., & Gunaratne, L. H. P. (2009). Chemical composition, biological effects of tannin and in vitro nutritive value of selected browse species grown in the West Australian Mediterranean environment. *Animal Feed Science and Technology*, 153(3–4), 203–215. <https://doi.org/10.1016/j.anifeedsci.2009.06.014>
- Lee, C., Araujo, R. C., Koenig, K. M., & Beauchemin, K. A. (2015). Effects of encapsulated nitrate on eating behavior, rumen fermentation, and blood profile of beef heifers fed restrictively or ad libitum. *Journal of Animal Science*, 93(5), 2405–2418. <https://doi.org/10.2527/jas2014-8851>
- León-Castro, Y.; Olivares-Pérez, J.; Rojas-Hernández, S.; Villa-Mancera, A.; Valencia-Almazán, M.T.; Hernández-Castro, E.; Córdova-Izquierdo, A.; Jiménez-Guillén, R. (2015). Effect of three fodder trees on *Haemonchus contortus* control and weight variations in kids. *Ecosistemas Y Recursos Agropecuarios*, 2(5), 193–201.
- Lorenz, M. M., Alkhafadji, L., Stringano, E., Nilsson, S., Mueller-Harvey, I., & Udén, P. (2014). Relationship between condensed tannin structures and their ability to precipitate feed proteins in the rumen. *Journal of the Science of Food and Agriculture*, 94(5), 963–968. <https://doi.org/10.1002/jsfa.6344>
- Nutritional Research Council (NRC). (2007). *Nutrient requirements of small ruminants*. Washington D.C. USA: National Academies Press. Retrieved from <https://www.nap.edu/read/11654/chapter/1>
- Olivares-Pérez, J., Avilés-Nova, F., Albarrán-Portillo, B., Castelan-Ortega, O. A., & Rojas-Hernández, S. (2013). Nutritional quality of *Pithecellobium dulce* and *Acacia cochliacantha* fruits, and its evaluation in goats. *Livestock Science*, 154(1–3), 74–81. <https://doi.org/10.1016/j.livsci.2013.02.017>
- Olivares-Pérez, J., Rojas-Hernández, S., Avilés-Nova, F., Camacho-Díaz, L. M., Cipriano-Salazar, M., Jiménez-Guillén, R., & Quiroz-Cardozo, F. (2016). USES OF NON-LEGUMINOUS TREES IN SILVOPASTORAL SYSTEMS IN THE SOUTH OF THE STATE OF MEXICO. *Ecosistemas Y Recursos Agropecuarios*, 3(8), 193–202.
- Olivares Perez, J., Aviles Nova, F., Albarran Portillo, B., Castelan Ortega, O. A., & Rojas Hernandez, S. (2013). Use of three fodder trees in the feeding of goats in the subhumid tropics in Mexico. *Tropical Animal Health and Production*, 45(3), 821–8. <https://doi.org/10.1007/s11250-012-0294-y>
- Ondiek, J. O., Ogore, P. B., Shakala, E. K., & Kaburu, G. M. (2013). Feed intake, digestibility and performance of growing small East African goats offered maize (*Zea mays*) stover supplemented with *Balanites aegyptiaca* and *Acacia tortilis* leaf forages. *Basic Research Journal of Agricultural Science and Review*, 2(1), 21–26.
- Oni, A. O., Onwuka, C. F. I., Oduguwa, O. O., Onifade, O. S., & Arigbede, O. M. (2008). Utilization of citrus pulp based diets and *Enterolobium cyclocarpum* (JACQ. GRISEB) foliage by West African dwarf goats. *Livestock Science*, 117(2–3), 184–191. <https://doi.org/10.1016/j.livsci.2007.12.010>
- Paengkoum, P. (2010). Effects of Neem (*Azadirachta indica*) and *Leucaena* (*Leucaena leucocephala*) Fodders on Digestibility, Rumen Fermentation and Nitrogen Balance of Goats Fed Corn Silage. *Journal of Animal and Veterinary Advances*, 9(5), 883–886.

- <https://doi.org/10.3923/javaa.2010.883.886>
- Paengkoum, P., & Paengkoum, S. (2010). ORIGINAL ARTICLE: Effects of supplementing rice straw with *Leucaena* (*Leucaena leucocephala*) and Madras thorn (*Pithecellobium dulce*) foliages on digestibility, microbial N supply and nitrogen balance of growing goats. *Journal of Animal Physiology and Animal Nutrition*, 94(5), e59–e65. <https://doi.org/10.1111/j.1439-0396.2009.00978.x>
- Patra, A. K., & Saxena, J. (2011). Exploitation of dietary tannins to improve rumen metabolism and ruminant nutrition. *Journal of the Science of Food and Agriculture*, 91(1), 24–37. <https://doi.org/10.1002/jsfa.4152>
- Pérez-Sato, M., Guerra-Medina, C. E., Becerril-Herrera, M., Soní-Guillermo, E., Ruiz-Sesma, B., Mendoza-Nazar, P., & Cobos-Peralta, M. A. (2011). Effect of two levels of tepozan bush (*buddleia cordata*) hay on the microorganisms in the rumen and the productive behavior in goats. *Tropical and Subtropical Agroecosystems*, 14(1), 129–135.
- Reis, G. L., Lana, A. M. Q., Mauricio, R. M., Lana, R. M. Q., Machado, R. M., Borges, I., & Neto, T. Q. (2010). Influence of trees on soil nutrient pools in a silvopastoral system in the Brazilian Savannah. *Plant and Soil*, 329(1), 185–193. <https://doi.org/10.1007/s11104-009-0144-5>
- Rojas-Hernández, S., Olivares-Pérez, J., Quiroz-Cardoso, F., Villa-Mancera, A., Cipriano Salazar, M., Camacho Díaz, L. M., R. P. A. (2016). Diagnosis of the palatability of fruits of three fodder trees in ruminants. *Ecosistemas y Recursos Agropecuarios*, 3(7), 121–127. <https://doi.org/http://natres.skc.rmuti.ac.th/pdf/Journal.PDF>
- Rojas-Hernandez, S., Olivares-Perez, J., Aviles-Nova, F., Villa-Mancera, A., Reynoso-Palomar, A., & Camacho-Díaz, L. M. (2015). Productive response of lambs fed *Crescentia alata* and *Guazuma ulmifolia* fruits in a tropical region of Mexico. *Tropical Animal Health and Production*, 47(7), 1431–1436. <https://doi.org/10.1007/s11250-015-0874-8>
- Rojas Hernández, S., Olivares Pérez, J., Elghandour, M. M. M. Y., Cipriano-Salazar, M., Ávila-Morales, B., Camacho-Díaz, L. M., ... A Cerrillo Soto, M. (2015). Effect of polyethylene glycol on in vitro gas production of some non-leguminous forage trees in tropical region of the south of Mexico. *Agroforestry Systems*, 89(4), 735–742. <https://doi.org/10.1007/s10457-015-9796-8>
- Sanon, H. O., Kaboré-Zoungana, C., & Ledin, I. (2008). Nutritive value and voluntary feed intake by goats of three browse fodder species in the Sahelian zone of West Africa. *Animal Feed Science and Technology*, 144(1–2), 97–110. <https://doi.org/10.1016/j.anifeedsci.2007.10.004>
- SAS. (2002). Statistical Analysis System: Guide for Personal Computers Version. NC, USA: Institute Inc. Cary.
- Steel, R. G. D., & Torrie, J. H. (2006). *Bioestadística: Principios y Procedimientos* (2nd ed.). México: McGraw-Hill/Interamericana de México.
- Sumano, H., & Ocampo, L. (2006). *Farmacología Veterinaria* (3rd ed.). Mc Graw Hill Interamericana. <https://doi.org/10.1017/CBO9781107415324.004>
- Supreena S. and Peangkoum P. (2011). Effect of *Leucaena leucocephala* (Lam.) and Sodium Lauryl Sulfate in Meat Goat Diets on Nematode Eggs and Protozoa Interaction in the Rumen. *Journal of Agricultural Science and Technology*, A(1), 1291–1294. Retrieved from <http://natres.skc.rmuti.ac.th/pdf/Journal.PDF>
- Táلامo, A., Barchuk, A., Cardozo, S., Trucco, C., Marás, G., & Trigo, C. (2015). Direct versus indirect facilitation (herbivore mediated) among woody plants in a semiarid Chaco forest: A spatial association approach. *Austral Ecology*, 40(5), 573–580. <https://doi.org/10.1111/aec.12224>
- Tamir, B., & Asefa, G. (2009). Effects of different forms of *Acacia saligna* leaves inclusion on feed intake, digestibility and body weight gain in lambs fed grass hay basal diet. *Animal Feed Science and Technology*, 153(1–2), 39–47. <https://doi.org/10.1016/j.anifeedsci.2009.05.010>
- Thang, C. M., Ledin, I., & Bertilsson, J. (2009). Effect of feeding cassava and/or *Stylosanthes* foliage on the performance of crossbred growing cattle. *Tropical Animal Health and Production*, 42(1), 1–11. <https://doi.org/10.1007/s11250-009-9378-8>

Recibido:
8/diciembre/2017

Aceptado:
26/octubre/2018

- Van Soest, P. J., Robertson, J. B., & Lewis, B. . (1991). Methods for Dietary Fiber, Neutral Detergent Fiber, and Nonstarch Polysaccharides in Relation to Animal Nutrition. *Journal of Dairy Science*, 74(10), 3583–3597. [https://doi.org/10.3168/jds.S0022-0302\(91\)78551-2](https://doi.org/10.3168/jds.S0022-0302(91)78551-2)
- Wang, Y. H., Xu, M., Wang, F. N., Yu, Z. P., Yao, J. H., Zan, L. S., & Yang, F. X. (2009). Effect of dietary starch on rumen and small intestine morphology and digesta pH in goats. *Livestock Science*, 122(1), 48–52. <https://doi.org/10.1016/j.livsci.2008.07.024>
- Williams, A. R., Ropiak, H. M., Fryganas, C., Desrues, O., Mueller-Harvey, I., & Thamsborg, S. M. (2014). Assessment of the anthelmintic activity of medicinal plant extracts and purified condensed tannins against free-living and parasitic stages of *Oesophagostomum dentatum*. *Parasites & Vectors*, 7(1), 518. <https://doi.org/10.1186/s13071-014-0518-2>
- Yinnesu, A., & Nurfeta, A. (2012). Effects of supplementing *Erythrina brucei* leaf as a substitute for cotton seed meal on growth performance and carcass characteristics of Sidama goats fed basal diet of natural grass hay. *Tropical Animal Health and Production*, 44(3), 445–451. <https://doi.org/10.1007/s11250-011-9916-z>